

Framingham Pediatrics

Routine Immunization and Screening Schedule

| | |
|-----------------|--|
| Newborn | HepB #1 |
| 2 Week | |
| 1 Month | |
| 2 Month | <u>Vaxelis</u> (DTaP #1, HepB #2, Hib #1,IPV #1) PCV20 #1, Rotavirus #1 |
| 4 Month | <u>Vaxelis</u> (DTaP #2, HepB #3, Hib #2, IPV #2) PCV20 #2, Rotavirus #2 |
| 6 Month | <u>Vaxelis</u> (DTaP #3, HepB #4, Hib #3, IPV #3) PCV20 #3, Rotavirus #3 (The 6 month visit must be <u>on or after</u> 6 months of age) |
| 9 Month | CBC & Lead screening |
| 12 Month | MMR #1, VARIVAX #1, PCV20 #4 SPOT vision screening (The 12 month visit must be <u>AFTER</u> the first birthday) |
| 15 Month | Pentacel (Dtap#4, Hib #4, IPV#4) Hep A #1 |
| 18 Month | |
| 24 Month | Hep A #2 CBC & Lead screening, SPOT vision screening (The 24 month visit must be <u>AFTER</u> the second birthday) |
| 2 ½ Year | At the discretion of the physician |
| 3 Year | CBC & Lead screening Begin yearly vision screening and BP check |
| 4 Year | Hearing Screening |
| 5 Year | <u>Kinrix</u> (DTaP #5, IPV #5), <u>Proquad</u> (MMR #2, VARIVAX #2) |

Yearly physical exams are recommended after 5 years of age.

| | |
|----------------|--|
| 9 Year | CBC, Cholesterol screening |
| 11 year | Tdap#1, HPV #1, MCV4 #1 HPV #2 given 6 months later or following year |
| 16 year | MCV4 #2 |
| 17 year | CBC, Cholesterol screening |

Entrance to 7th Grade requires completion of the Hepatitis B Series, Tdap Booster, documentation of chicken pox disease or Varivax, and 2 MMR vaccines.

MCV4 Booster - Given at age **16** if primary dose is administered at age 11-12yrs, Age 16-18yrs if primary dose is administered at 13-15yrs. **Booster not required if primary dose is administered after 16yrs.**

HPV vaccine if started on or after age 15: 3 dose series (0, 1-2 months, 6 months)

Minimal interval between dose 1 and 2 is 4 weeks, dose 2 and 3 is 12 weeks and dose 1 and 3 is 5 months

Earliest age for 1st dose of **Vaxelis** (DTaP, Hep B, IPV Hib) and PCV15 is 6 weeks old.

1st dose of **Rotavirus** vaccine must be given prior to 12 weeks of age, last dose prior to 32 weeks

Minimal interval between vaccines **Vaxelis** (DTaP, Hep B, IPV, Hib) and PCV15 is 4 weeks.

Hepatitis B Vaccine for older patients requires minimum 4 week interval between 1st and 2nd doses with 3rd dose given 6-12 months later (minimum intervals of 4 months between 1st and 3rd doses and 2 months between 2nd and 3rd doses).